



## Does secure base script knowledge mediate associations between observed parental caregiving during childhood and adult romantic relationship quality and health?

Or Dagan, Renate S. M. Buisman, Marissa D. Nivison, Theodore E. A. Waters, Brian E. Vaughn, Kelly K. Bost, Maria E. Bleil, Deborah Lowe Vandell, Cathryn Booth-LaForce & Glenn I. Roisman

To cite this article: Or Dagan, Renate S. M. Buisman, Marissa D. Nivison, Theodore E. A. Waters, Brian E. Vaughn, Kelly K. Bost, Maria E. Bleil, Deborah Lowe Vandell, Cathryn Booth-LaForce & Glenn I. Roisman (2021) Does secure base script knowledge mediate associations between observed parental caregiving during childhood and adult romantic relationship quality and health?, Attachment & Human Development, 23:5, 643-664, DOI: [10.1080/14616734.2020.1836858](https://doi.org/10.1080/14616734.2020.1836858)

To link to this article: <https://doi.org/10.1080/14616734.2020.1836858>



Published online: 27 Oct 2020.



Submit your article to this journal [↗](#)



Article views: 184



View related articles [↗](#)



View Crossmark data [↗](#)

ARTICLE



## Does secure base script knowledge mediate associations between observed parental caregiving during childhood and adult romantic relationship quality and health?

Or Dagan<sup>a</sup>, Renate S. M. Buisman<sup>b</sup>, Marissa D. Nivison<sup>c</sup>, Theodore E. A. Waters<sup>d</sup>, Brian E. Vaughn<sup>e</sup>, Kelly K. Bost<sup>f</sup>, Maria E. Bleil<sup>g</sup>, Deborah Lowe Vandell<sup>h</sup>, Cathryn Booth-LaForce<sup>i</sup> and Glenn I. Roisman<sup>c</sup>

<sup>a</sup>Department of Psychology, Stony Brook University, New York, NY, USA; <sup>b</sup>Vrije Universiteit Amsterdam, Amsterdam, The Netherlands; <sup>c</sup>Institute of Child Development, University of Minnesota, Minnesota, MN, USA; <sup>d</sup>Department of Psychology, New York University, Abu-Dhabi, United Arab Emirates; <sup>e</sup>Department of Human Development and Family Studies, Auburn University, Auburn, AL, USA; <sup>f</sup>Human and Community Development, University of Illinois, Champaign, IL, USA; <sup>g</sup>School of Nursing, University of Washington, Seattle, WA, USA; <sup>h</sup>School of Education, University of California, Irvine, CA, USA; <sup>i</sup>University of Washington, Seattle, WA, USA

### ABSTRACT

Increasingly, attachment representations are being assessed via *secure base script knowledge* – the degree to which individuals show awareness of the temporal-causal schema that summarizes the basic features of seeking and receiving effective support from caregivers during times of need. Limited research has assessed the links between secure base script knowledge and aspects of adult functioning and the role that secure base script knowledge may play in accounting for associations between early caregiving quality and adulthood functioning. We used follow-up assessments of the NICHD Study of Early Child Care and Youth Development cohort ( $N = 585$ ) to examine whether secure base script knowledge at age 18 years: (a) is associated with later romantic relationship quality, depressive symptoms, and body mass index (BMI) at age 26 years, and (b) mediates expected associations between the quality of maternal and paternal sensitivity across the first 15 years of life and age-26 outcomes. More access to, and elaborated knowledge of the secure base script predicted less extreme hostility with romantic partners, and better emotional and physical health. Moreover, secure base script knowledge mediated the links between early maternal and paternal sensitivity and both later romantic partner hostility and depressive symptoms, but not BMI.

### KEYWORDS

Parental sensitivity; secure base script knowledge; romantic relationships; depressive symptoms; physical health

Attachment theory posits that the quality of early care, reflected in the caregiver's provision of contingent, appropriate, and consistent (i.e., sensitive) responses to a child's emotional needs, supports the development and consolidation of mental representations of the attachment figures' availability in times of distress (Ainsworth et al., 1978; Bretherton & Munholland, 2008; Marvin & Britner, 2008). The literature largely

**CONTACT** Or Dagan  [or.dagan@stonybrook.edu](mailto:or.dagan@stonybrook.edu)  Department of Psychology, Stony Brook University; Glenn I. Roisman  [roism001@umn.edu](mailto:roism001@umn.edu)  Institute of Child Development, University of Minnesota

supports the claim that such attachment representations, in turn, contribute to individual differences in adult functioning, including interpersonal functioning with romantic partners (Collins & Sroufe, 1999), emotional well-being (Dagan, Asok et al., 2018a, 2020; Stovall-McClough & Dozier, 2016), and even aspects of physical health (Dagan, Facompré et al., 2018b; Ehrlich & Cassidy, 2019; Farrell et al., 2019).

Research suggests that attachment representations can be conceptualized, in part, as a cognitive script summarizing the quality of early caregiving interactions (e.g., Bretherton, 1985, 1987) – specifically, a *secure base* script (Waters & Waters, 2006). The secure base script is a temporal-causal generalization, or schema, that summarizes the basic features of seeking/receiving effective support from an attachment figure when this support is needed. More specifically, a history of reliable and effective secure base support is assumed to become generalized as an expectation that primary caregivers (and, indeed, secure base figures in general) will consistently be available, recognize and promptly respond to attachment-related threats, and will possess both the wisdom and the competence to restore balance to ongoing activities (Waters & Facompré, [in press](#)).

### Secure base script knowledge and adult outcomes

Given that secure base script knowledge assesses individual's expectations of close others' availability and support during times of distress, it is plausible to expect such representations influence one's romantic relationship quality, mood, and health status. However, limited research has been conducted regarding the links between secure base script knowledge and aspects of adjustment in the adult years. Instead, in many investigations to date, secure base script knowledge has been studied either in relation to its caregiving antecedents (Nivison et al., 2020; Schoenmaker et al., 2015; Steele et al., 2014; Vaughn et al., 2016; Waters et al., 2017) or its role in shaping both adults' caregiving and attachment security in the next generation (Monteiro et al., 2008; Vaughn et al., 2007; Waters et al., 2018; see Waters & Roisman, 2019, for a review). Nonetheless, some research regarding the predictive significance of secure base script knowledge for adult functioning has been conducted, with a focus on three developmentally salient indicators of individual adjustment in the adult years: interpersonal functioning with adult romantic partners, emotional well-being, and physical health.

Two studies have thus far examined the relevance of secure base script knowledge for the quality of adults' romantic relationships. Waters et al. (2013) found that secure base script knowledge predicted care-seeking behavior with a romantic partner during a lab based conflict discussion task. Subsequently, Waters et al. (2018) replicated and extended these findings, reporting significant bivariate associations between secure base script knowledge and the quality of interaction with romantic partners, self-reported relationship satisfaction, and an interview-based assessment of effectiveness in romantic relationships in the higher-risk sample of the Minnesota Longitudinal Study of Risk and Adaptation (MLSRA; Sroufe et al., 2005).

Other studies have suggested a link between the quality of romantic relationships and attachment representations assessed via the coherence of discourse during the Adult Attachment Interview (AAI; Main et al., 2003–2008) which is moderately correlated with secure base script knowledge (e.g. Steele et al., 2014). In short, higher coherence scores are coded in interviews with discourse which entails descriptions of experiences and

relationships with childhood caregivers in an internally consistent but not emotionally overwrought manner (e.g., Roisman, 2009). Secure attachment representations (i.e. high coherence scores) were associated with behaviors that indicate care seeking and caregiving with romantic partners (Crowell et al., 2002), and prospectively predicted romantic relationship quality, assessed via observer-rated couple interactions and self-reported relationship quality (Holland & Roisman, 2010; Roisman et al., 2001).

Secure base script knowledge was also found to predict symptoms of psychopathology, including depressive symptoms, in middle childhood (Ruiz et al., 2019; Waters, Fraley et al., 2015). The link between secure base script knowledge and depressive symptoms (i.e., emotional health) in adults has yet to be assessed; however, a recent meta-analysis summarizing research regarding the association between attachment representations (assessed using AAI coherence) and depressive symptoms in over 4,000 adults (Dagan, Facompré et al., 2018b) reported that adults classified as insecurely attached had more depressive symptoms than did their secure counterparts. These results support the notion that an individual's mental representations regarding past relationships with primary caregivers are significantly related to their depressive symptomatology in adulthood.

Finally, building on accumulating evidence suggesting that the quality of early experiences with one's caregivers "gets under the skin" to influence adult physical health (e.g. Puig et al., 2013), one study assessed the predictive significance of secure base script knowledge for adult physical health. Farrell et al. (2019) found that secure base script knowledge at ages 19 and 26 years was negatively associated with a cardiometabolic index composed of four anthropometric indicators and biomarkers of health, including an objective assessment of body mass index (BMI), at ages 37 and 39 years. Other studies reporting on adult attachment representations as assessed in the AAI likewise support the link between secure base script knowledge and obesity. For example, one study that assessed 215 adults at midlife observed that individuals who exhibited lower coherence of discourse in the AAI had higher metabolic syndrome indicators, including obesity, elevated blood pressure, elevated fasting glucose, high triglycerides, and low high-density lipoprotein cholesterol (Davis et al., 2014) – all of which are associated with higher BMI (Rodriguez et al., 2015). Taken together, these studies and others (for a review, see Diener et al., 2016) support the claim that attachment representations in general, and secure base script knowledge in particular, may play a significant role in regulation of physiological stress response systems, which in turn has implications for BMI and other aspects of physical health (Lambert & Lambert, 2011).

### **Secure base script knowledge may account for associations between early sensitivity and adult outcomes**

Studies have shown that early parental sensitivity predicts higher quality of secure base script knowledge in later life (Schoenmaker et al., 2015; Steele et al., 2014; Vaughn et al., 2016; Waters et al., 2017). Moreover, secure base script knowledge is thought to influence adult functioning by constituting a heuristic for guiding behaviors (Schank & Abelson, 1977), especially during distress (Waters & Waters, 2006). Taken together, secure base script knowledge may play a role in *explaining* the association between parental sensitivity and adult functioning. Thus, we turn to evidence for: (a) the link between early

sensitive care and adult outcomes, and (b) the role of secure base script knowledge and its attachment representation correlates in mediating these associations.

### *From early sensitivity to adult outcomes*

Longitudinal studies of human development have reported consistent associations between the quality of early child–parent interactions and romantic functioning in young adulthood. For example, several reports based on the MLSRA (Englund et al., 2011; Raby, Lawler et al., 2015; Raby, Roisman et al., 2015) demonstrate that maternal sensitivity during infancy and childhood is reliably associated with adult romantic relationship competence; the latter was assessed via both interviews regarding the stability and quality of past and current romantic relationships and electrodermal activity during conflict discussions with their adult partners. Other studies (e.g., Conger et al., 2000) have supported the link between early child–parent interactions and behaviors in romantic relationships during adulthood, as assessed in observed conflict tasks with romantic partners and self-reported thoughts and feelings regarding their romantic relationships.

Maternal sensitivity during infancy and childhood has also been linked to later depressive symptoms. A longitudinal study found a link between maternal sensitivity and depressive symptoms in MLSRA participants between the ages 16 and 17.5 years (Duggal et al., 2001). Compared with individuals who experienced sensitive care in infancy and childhood, those who experienced more maternal *insensitive* care during infancy and childhood tended to report more depressive symptoms in adolescence.

Accumulating evidence has also demonstrated that the quality of early parent–child relationships is an important predictor of obesity later in life. For example, Anderson et al. (2012) found that lower maternal sensitivity observed in the first 3 years of life was associated with higher BMI at age 15 years. Independent bodies of research indicate robust negative associations between early maternal care and childhood obesity (for review, see Anderson & Keim, 2016). Moreover, childhood obesity and adulthood obesity are positively correlated (for review, see Simmonds et al., 2016). Taken together, these findings lend further support to our hypothesized link between early caregiving experiences and obesity later in life.

### *The mediating role of secure base script knowledge*

Evidence is limited for the role that attachment representations in general, and secure base script knowledge specifically, play in explaining the multiple links between parental sensitivity and adult outcomes. Using a sample of at-risk individuals assessed across time in the MLSRA, Roisman et al. (2001) found a significant association between child–mother dyadic behaviors observed at age 13 years and dyadic behaviors with romantic partners at age 20–21 years. Moreover, this link was mediated by attachment representations assessed at age 19 years, as inferred from their coherence of discourse scored from their AAI transcripts.

Recently, the role of secure base script knowledge was also examined as a mediator between early sensitivity and health outcomes in the MLSRA. Using MLSRA data, Farrell et al. (2019) found that maternal sensitivity in the first four years of life predicted scores on a cardiometabolic index, including BMI, 30 years later. Importantly, Farrell et al. (2019)

showed that attachment representations at age 19 and 26 years, as indicated by secure base script knowledge, partially mediated this link.

## The present study

Taken together, evidence suggests that attachment representations, as reflected in secure base script knowledge, partially account for associations between early experiences with primary caregivers and adult functioning in an at-risk sample (i.e., the MLSRA). In the current study, we sought to leverage the larger, normative-risk NICHD Study of Early Child Care and Youth Development (SECCYD) – a 26-year prospective, longitudinal investigation of human development – to attempt to replicate and extend the pattern of results obtained thus far. In addition, we aimed to extend previous findings in two ways. First, because studies have yet to assess the role secure base script knowledge may play in emotional health during adulthood, we incorporated depressive symptoms as an adult outcome. Second, the MLSRA did not acquire measures of paternal sensitivity, so previously reported studies could only assess the effects of *maternal* sensitivity on subsequent outcomes via secure base script knowledge. The SECCYD included assessments of sensitivity with both maternal and paternal caregivers so we were able to consider effects of each parent as a predictor of both secure base script knowledge and relevant relationship and health outcomes.

We hypothesized, first, that secure base script knowledge assessed at age 18 years would be associated with adult functioning, as indicated by self-reported romantic relationship quality (i.e. perceived romantic partner's warmth and hostility), depressive symptoms, and BMI (derived from self-reported height and weight) at age 26. Second, we tested whether secure base script knowledge mediated the hypothesized associations between the quality of early parental caregiving – as reflected in observations of maternal and paternal sensitivity across the first 15 years of life – and reported romantic relationship quality, depressive symptoms, and BMI in young adulthood, around age 26 years. Of note, little research exists on the differential effects of early paternal and maternal sensitivity on adult outcomes. Given that previous research indicated similar maternal and paternal sensitivity effects on secure base script knowledge in the current sample (Steele et al., 2014), we hypothesized that secure base script knowledge would mediate the expected links between both maternal and paternal sensitivity and adult outcomes.

## Method

### Participants

The SECCYD is an ongoing longitudinal study that has followed a sample of children and their families (original  $N = 1364$ ) at 10 sites across the United States from birth to age 15 years. In addition, follow-ups have occurred, most notably for this report at age 18 years and then again at age 26 years. The primary purpose of the original SECCYD was to examine the characteristics of relevant contexts (e.g. childcare, home, school, neighborhood) and individual characteristics of the child and the family in relation to children's developmental outcomes (e.g. socio-emotional, cognitive, language, health) (for detailed information, see <https://www.nichd.nih.gov/research/supported/seccyd>).

The analytic frame for this report was the subsample of young adults who had completed an Attachment Script Assessment at an age 18-year follow-up of the SECCYD ( $N = 673$ ; see Steele et al., 2014), and who also completed any of the five relevant outcome measures (i.e. three romantic relationship indices, depressive symptoms, and BMI) in an online survey at age 26 years. Accordingly, romantic relationship outcome analyses (i.e., partner's warmth, and partner's moderate and extreme hostility) included data from 400 participants (a subset of age 26-year-old participants who were also in a current romantic relationship). Depressive symptoms outcome analyses included data from 584 participants and BMI outcome analyses included data from 578 participants. Overall, 54.4% of participants were female, and the majority were White/non-Hispanic (80.7%) and middle-class (average of income-to-needs ratio was 4.22;  $SD = 2.89$ ). See Table 1 for additional sample characteristics.

### *Attrition analysis*

We compared demographic characteristics for the 26-year outcomes subsample (hereafter "26-year sample";  $n = 585$ ) with those in the original SECCYD sample not included in the current sample (hereafter "attriters";  $n = 779$ ). The 26-year sample comprised a higher percentage of White/non-Hispanic (80.7%) and females (54.4%) compared with the attriters (73.2% and 43.8%, respectively),  $\chi^2(1, N = 1,364) = 14.99, p < .001, d = .21$ , and  $\chi^2(1, N = 1,364) = 10.46, p = .001, d = .18$ , respectively. The 26-year sample participants had higher Full Scale IQ scores ( $M = 108.35, SD = 13.80$ ) compared with the attriters ( $M = 104.93, SD = 15.02$ ),  $t(1010) = 3.42, p < .001, d = 0.24$ . The 26-year sample participants also had mothers with significantly more years of education ( $M = 14.68, SD = 2.39$ ) compared with the attriters ( $M = 13.90, SD = 2.55$ ),  $t(1361) = 5.69, p < .001, d = 0.32$ ; and had higher family income-to-needs ratio ( $M = 4.18, SD = 2.86$ ) compared with the attriters ( $M = 3.40, SD = 3.07$ ),  $t(1354) = 4.73, p < .001, d = 0.26$ . As indicated by Cohen's  $d$  effect sizes, all such differences between the two samples were small in magnitude as per Cohen's (1992) criteria.

### *Measures*

#### *Parental sensitivity*

Maternal and paternal sensitivity were derived from direct observations of age-appropriate tasks performed by participants and their mothers/fathers; for information regarding the tasks and their scoring systems see Booth-LaForce et al. (2014), and Fraley et al. (2013). Observations took place at age 6, 15, 24, and 36 months with mothers only, and at age 54 months, Grades 1, 3, and 5, and age 15 years with mothers and fathers separately. Following prior studies using data from the SECCYD, separate observed maternal and paternal sensitivity composites were calculated by standardizing and averaging scores across time points. As noted in Steele et al. (2014), in the current sub-sample of the SECCYD Cronbach's alpha composite sensitivity scores were .83 and .72 (for maternal and paternal sensitivity, respectively).

#### *Secure base script knowledge*

The adolescent version of the Attachment Script Assessment (ASA; Dykas et al., 2006; Waters & Waters, 2006) was used to assess participants' secure base script knowledge. This

**Table 1.** Correlations among parental sensitivity, secure base script knowledge, and adult functioning at age 26 years.

Variable	1	2	3	4	5	6	7	8	9	10	11	12	13
1. SBSK <sup>1</sup>	-												
2. Partner Warmth <sup>2</sup>	.03	-											
3. Partner MH <sup>2</sup>	-.10	-.54**	-										
4. Partner EH <sup>2</sup>	-.18**	-.32**	.38**	-									
5. Depression <sup>2</sup>	-.13**	-.29**	.31**	.19**	-								
6. BMI <sup>2</sup>	-.10*	-.08	.15**	.04	.07	-							
7. Maternal sensitivity	.28**	.08	-.12**	-.17**	-.09*	-.25**	-						
8. Paternal sensitivity	.29**	.07	-.14**	-.14**	-.08	-.21**	.42**	-					
9. Child gender	.20**	.02	.08	-.12*	.05	-.05	.11**	.08	-				
10. Child ethnicity	.13**	.09	-.03	.13**	-.12**	-.04	.36**	.09*	.02	-			
11. Maternal education	.19**	-.02	-.04	-.03	-.08	-.19**	.49**	.22**	.06	.25**	-		
12. Income/needs ratio	.18**	-.00	-.13*	-.08*	-.15**	-.25**	.43**	.24**	.05	.25**	.56**	-	
13. IQ	.21**	.00	-.09	-.08	-.01	-.21**	.45**	.27**	-.02	.27**	.46**	.38**	-
Means	3.78	4.35	1.76	1.05	1.75	26.46	0.00	0.00	54.4 <sup>a</sup>	80.7 <sup>b</sup>	14.68	4.22	-
SDs	1.05	0.63	0.63	0.22	0.60	6.19	1.00	1.00	-	-	2.39	2.89	-

<sup>1</sup>Secure Base Script Knowledge at Age 18; <sup>2</sup>At Age 26; <sup>a</sup> Percent female; <sup>b</sup> Percent White/Non-Hispanic; MH = Moderate Hostility; EH = Extreme Hostility; BMI = Body Mass Index. Child gender (1 = male, 2 = female); Child ethnicity (0 = other, 1 = White, non-Hispanic); Income/needs ratio = income-to-needs-ratio. \*  $p < .05$ . \*\*  $p < .01$ .

version of the ASA consisted of four stories, with a 12 word-prompt outline for each story; two mother-related stories (“The Party,” and either “Acne” [completed by girls] or “The Haircut” [completed by boys]), and two father-related stories (“The Tennis Match” and “Studying for an Exam”). Participants first completed a practice story (“A Trip to the Beach”; not coded) to ensure they understood the task. Participants were asked to tell the “best story” they could tell using the outline of each story. The task was not timed, and participants were asked to tell a story of approximately one page in length. Audio recordings of the ASA session were transcribed verbatim, and each story was coded on a 7-point scale, from 1 (no secure base script content) to 7 (extensive secure base script organization with substantial elaboration of script relevant details). Following the convention in prior studies using the ASA (e.g., ; Vaughn et al., 2006) and evidence that the scores share loadings on a single factor, a composite secure base script knowledge score was calculated by averaging the ASA scores across the four stories (for elaborated reliability data see Steele et al., 2014). All ASA stories were double coded, and intraclass correlation coefficient (ICC) was .95. Cronbach’s alpha for the four-story composite of the ASA sub-sample of the SECCYD was .78.

### *Romantic relationship quality*

At age 26 years, romantic relationship quality was assessed with the Behavioral Affect Rating Scale (BARS; Conger, 1989). The BARS measures warmth and hostility within a close relationship and was developed for research with rural families in Iowa. It has been shown to be a reliable and valid measure of warmth and hostility within family relationships across multiple ethnic groups (e.g. Taylor et al., 2012).

In this study, participants reported on their romantic partners’ behavior towards them. The 20-item version was used, yielding scores for three subscales: Warmth, Moderate Hostility, and Extreme Hostility. Response categories were modified from the original 7-point Likert-type scale used in Conger’s Iowa sample and the 4-point rating scale used in Conger’s Mexican-American sample to a 5-point Likert-type scale that ranged from 1 (*never*) to 5 (*always*) used by Eccles and colleagues in the age-26 follow-up (Wave 9) of the Michigan Study of Achievement and Life Transitions. Sample items include: “Last month, how often did your partner listen carefully to your point of view?” (Warmth); “Last month, how often did your partner criticize you or your ideas?” (Moderate Hostility); and “Last month, how often did your partner throw something at you?” (Extreme Hostility). Higher scores on each subscale indicated higher levels of warmth, moderate and extreme hostility, respectively.

In the current study, internal consistency of both romantic partner’s Warmth and Moderate Hostility composite scores were high ( $\alpha = .90$  and  $.84$ , respectively), and the reliability of the romantic partner’s Extreme Hostility composite score was lower (as expected given that only three indices were assessed;  $\alpha = .60$ ). We elected to use the standard composite for romantic partner’s Extreme Hostility (i.e. a mean of three items that assess behaviors of hit/push/grab/shove person, throw something at person, and threaten person with a knife/gun) because the behaviors referenced are low base rate in the population and are *prima facie* indicators of extreme hostility. In the current sample, the romantic partner’s Extreme Hostility composite scores were non-parametrically distributed (skewness of 6.17;  $SE = 0.12$ ), with 17% of the subsample’s participants endorsing the presence of any of the potential partner’s extreme hostility indices.

### *Depressive symptoms*

The Center for Epidemiological Studies-Depression (CES-D; Radloff, 1977) scale was used to measure participants' depressive symptoms within the last week. The 20-item measure consists of depressive symptoms, such as restless sleep, poor appetite, and feeling lonely. Response options range from 1 (rarely or none of the time) to 4 (most or almost all of the time). Items that indicated positive feelings (i.e. "I felt hopeful about the future") were reverse coded such that higher scores indicated greater depressive symptoms. The CES-D has proven to be a reliable and valid measure of the major facets of depression (e.g. Cosco et al., 2017), and its internal consistency in the current study was high ( $\alpha = .94$ ).

### *BMI*

Participants' age-26 body mass index (BMI) was calculated based on their self-reported height (in feet) and weight (in pounds). Self-report and objective measures of height, weight, and overall BMI were shown to be very highly correlated (e.g. Wilson et al., 2019). Height and weight were first converted to meters and kilograms, respectively. Next, weight was divided by height squared to yield BMI.

### *Covariates*

Covariates included child ethnicity (1 = White/non-Hispanic, 0 = Other), child gender (1 = Male, 2 = Female), child's IQ, maternal years of education, and family income. Child's IQ was quantified as the full-scale IQ obtained in the Wechsler Abbreviated Scale of Intelligence (1999) in fourth grade. Family income was operationalized by averaging the income-to-needs ratio (total family income divided by the year-specific poverty threshold for the appropriate family size) across the first 15 years of the individual's life. Income-to-needs ratio was calculated separately at 1, 6, 15, 24, 36, and 54 months; during Grade 1, 3, 4, 5, and 6; and at age 15 years.

### *Analytic strategy*

First, Pearson correlations were computed to examine associations among the study variables. Next, to examine whether secure base script knowledge-mediated associations between both maternal and paternal sensitivity across the first 15 years of life and adult functioning at age 26 years, we conducted mediation analyses using the PROCESS macro for SPSS, version 3.4 (Hayes, 2017; available at <http://processmacro.org/index.html>). Using a percentile bootstrap estimation method (Preacher & Hayes, 2008), PROCESS macro allows for non-parametric distributions. Given the low base rate of partner's extreme hostility indices, we also ran a binary logistic mediation analysis to assess this outcome dichotomously (i.e., whether participants endorsed none or any extreme partner hostility indices).

Consistent with the approach of Farrell et al. (2019), a separate model was conducted for each outcome measure, and each model was independently assessed with both maternal and paternal sensitivity as independent variables, as well as with and without covariates (child ethnicity, child gender, maternal years of education, and family income). Indirect effects were tested using a percentile bootstrap estimation approach with 10,000 samples, as recommended by Hayes (2017). When significant, indirect effect sizes were calculated based on Preacher and Kelley (2011) kappa-squared ( $\kappa^2$ ) effect size metric

recommendation for mediation models, with .01, .09, and .25 representing small, medium, and large effect sizes, respectively. Finally, we probed significant mediation models further to assess whether one caregiver's sensitivity (e.g., maternal sensitivity) affected the outcome of interest above and beyond the other caregiver's sensitivity (e.g., paternal sensitivity). To this end, we re-ran significant mediation analyses with one parental sensitivity score as an independent variable and the other parental sensitivity score as a covariate.

## Results

### *Correlational analyses*

Table 1 reports correlations among all study variables, derived from the sub-samples described above. Secure base script knowledge was negatively associated with romantic partner extreme hostility ( $r = -.18, p < .001$ ), depressive symptoms ( $r = -.13, p = .002$ ), and BMI ( $r = -.10, p = .02$ ). These correlations indicate that more access to, and elaborated knowledge of the secure base script at 18 years was associated with less perceived extreme hostility from a romantic partner, fewer depressive symptoms, and a lower BMI at 26 years of age. The secure base script score was not significantly associated with either partner warmth ( $r = .03, p = .62$ ) or partner moderate hostility ( $r = .18, p = .054$ ).

Consistent with Steele et al. (2014), both maternal ( $r = .28, p < .001$ ) and paternal ( $r = .29, p < .001$ ) sensitivity were positively correlated with secure base script knowledge, suggesting that higher parental sensitivity across the first 15 years of life is associated with more secure base script knowledge at age 18 years. In addition, both maternal and paternal sensitivity were significantly and negatively correlated with both partner moderate hostility ( $r = -.12, p = .02$  and  $r = -.14, p = .007$ , respectively) and extreme hostility ( $r = -.17, p = .001$  and  $r = -.14, p = .006$ , respectively). Maternal ( $r = -.09, p = .02$ ), but not paternal ( $r = -.08, p = .09$ ) sensitivity was negatively correlated with depressive symptoms, indicating that more sensitivity received from mother across the first 15 years of life was associated with fewer self-reported depressive symptoms at age 26 years. Finally, both maternal ( $r = -.25, p < .001$ ) and paternal ( $r = -.21, p < .001$ ) sensitivity were associated with lower BMI at age 26 years.

### *Mediational analyses*

We analyzed all hypothesized models using ordinary least squares regression analysis, with and without covariates. We found that results were highly similar in these cases (see Tables 2 and 3 for results of both types of analyses). For the sake of simplicity, the results below are reported only on the models that were adjusted for covariates (i.e., child gender, age, and IQ, maternal education, and income/needs ratio).

### *Romantic relationship quality*

We tested the hypothesis that secure base script knowledge mediates the link between parental sensitivity across the first 15 years and romantic relationship quality, as assessed via three self-reported indicators at age 26 years: partner warmth, moderate hostility, and extreme hostility.

With respect to partner's warmth, results indicated that both maternal ( $b = .20, SE = .06, p < .001$ ) and paternal ( $b = .20, SE = .05, p < .001$ ) sensitivity were significant predictors of secure base script knowledge. However, when adjusted for parental sensitivity, secure base script knowledge did not significantly predict partner's warmth ( $b = .00, SE = .03, p = .99$  and  $b = .01, SE = .04, p = .94$ , respectively). These results did not support our mediational hypothesis.

A similar pattern of results was obtained when we examined partner's moderate hostility as the dependent variable. The indirect link between parental sensitivity and partner's moderate hostility through secure base script knowledge was not significant. Although both maternal ( $b = .20, SE = .06, p < .001$ ) and paternal ( $b = .20, SE = .05, p < .001$ ) sensitivity were significant predictors of secure base script knowledge, in both models secure base script knowledge was not a significant predictor of partner's moderate hostility when adjusting for parental sensitivity ( $b = -.03, SE = .03, p = .44$  and  $b = -.02, SE = .04, p = .67$ , respectively). Again, the results were not consistent with the mediational hypothesis.

In contrast, when we modeled partner extreme hostility, results indicated that both maternal ( $b = .20, SE = .06, p < .001$ ) and paternal ( $b = .20, SE = .05, p < .001$ ) sensitivity were significant predictors of secure base script knowledge, which in turn significantly predicted partner's extreme hostility ( $b = -.03, SE = .01, p < .03$  and  $b = -.02, SE = .00, p = .04$ , respectively). Moreover, coefficients for the indirect effects in both cases were significant, as evidenced by the absence of a zero value within the bootstrap confidence intervals ( $b = -.01, 95\% CI = -.02, -.01$  and  $b = -.01, 95\% CI = -.01, -.001$ , respectively). These results support the mediational hypothesis. Approximately 5% of the variance in partner's extreme hostility was accounted for by the predictors in both models ( $R^2 = .05$ ).

**Table 2.** Path coefficients and indirect effects for mediation models with **maternal sensitivity** as independent variable and secure base script knowledge as mediator.

Outcome Variable (n)	Path A			Path B			Path C'			Indirect Effects		
	b	SE	t	b	SE	t	b	SE	t	Effect	BootLLCI	BootULCI
Partner Warmth												
No covariates (400)	.32**	.05	6.33	.00	.03	.01	.05	.04	1.52	.00	-.02	.02
With covariates (390)	.20**	.06	3.30	.00	.03	-.01	.05	.04	1.17	.00	-.01	.01
Partner Moderate Hostility												
No covariates (400)	.32**	.05	6.33	-.04	.03	-1.27	-.07	.03	-1.89	-.01	-.03	.01
With covariates (390)	.20**	.06	3.30	-.03	.03	-.88	-.03	.04	-.77	-.01	-.02	.01
Partner Extreme Hostility												
No covariates (400)	.32**	.05	6.33	-.03**	.01	-2.83	-.03*	.01	-2.47	-.01	-.02	-.01
With covariates (390)	.20**	.06	3.30	-.03*	.01	-2.20	-.02	.01	-1.62	-.01	-.02	-.01
Depressive Symptoms												
No covariates (584)	.29**	.04	6.85	-.07*	.03	-2.57	-.04	.08	-1.47	-.02	-.03	-.01
With covariates (570)	.18**	.05	3.59	-.07**	.03	-2.70	.01	.03	-.03	-.01	-.03	-.001
BMI												
No covariates (578)	.29**	.04	6.67	-.18	.26	-.72	-1.62**	.28	-5.88	-.05	-.22	.11
With covariates (564)	.18**	.05	3.49	-.15	.25	-.58	-.92**	.31	-2.95	-.03	-.14	.08

Path A = Direct effect of the independent variable (Maternal Sensitivity) on the mediator (Secure Base Script Knowledge; SBSK); Path B = Direct effect of the mediator (SBSK) on the dependent variable; Path C' = Direct effect of the independent variable (Paternal Sensitivity) on the dependent variable, controlling for the mediator (SBSK); BootLLCI = Bootstrapping lower limit confidence interval; BootULCI = Bootstrapping upper limit confidence interval; SE = Standard Error; BMI = Body Mass Index. Covariates are Child Gender, Child Ethnicity, Maternal Education, Income-to-Needs Ratio, and Full Scale IQ in 4<sup>th</sup> Grade. \*  $p < .05$ . \*\*  $p < .01$ .

**Table 3.** Path coefficients and indirect effects for mediation models with **paternal sensitivity** as independent variable and secure base script knowledge as mediator.

Outcome Variable (n)	Path A			Path B			Path C'			Indirect Effects		
	<i>b</i>	<i>SE</i>	<i>t</i>	<i>b</i>	<i>SE</i>	<i>t</i>	<i>b</i>	<i>SE</i>	<i>t</i>	<i>Effect</i>	<i>BootLLCI</i>	<i>BootULCI</i>
<b>Partner Warmth</b>												
No covariates (357)	.29**	.05	6.12	.01	.03	.37	.04	.03	1.20	.00	-.02	.02
With covariates (348)	.20**	.05	4.09	.00	.04	.08	.04	.03	1.23	.00	-.01	.01
<b>Partner Moderate Hostility</b>												
No covariates (357)	.29**	.05	6.12	-.04	.04	-1.07	-.07*	.03	-2.24	-.01	-.03	.01
With covariates (348)	.20**	.05	4.09	-.02	.04	-.43	-.05	.03	-1.45	.00	-.02	.01
<b>Partner Extreme Hostility</b>												
No covariates (357)	.29**	.05	6.12	-.03**	.01	-2.89	-.02	.01	-1.75	-.01	-.02	-.01
With covariates (348)	.20**	.05	4.09	-.02*	.01	-2.08	-.01	.01	-1.27	-.01	-.01	-.001
<b>Depressive Symptoms</b>												
No covariates (510)	.28**	.04	6.79	-.07*	.03	-2.40	-.03	.03	-.96	-.02	-.04	-.01
With covariates (500)	.21**	.04	4.89	-.07*	.03	-2.34	-.02	.03	-.63	-.01	-.03	-.001
<b>BMI</b>												
No covariates (506)	.28**	.04	6.72	-.27	.28	-.95	-1.21**	.27	-4.43	-.07	-.25	.09
With covariates (496)	.21**	.04	4.80	-.05	.28	-.19	-.77*	.28	-2.80	-.01	-.14	.12

Path A = Direct effect of the independent variable (Paternal Sensitivity) on the mediator (Secure Base Script Knowledge; SBSK); Path B = Direct effect of the mediator (SBSK) on the dependent variable; Path C' = Direct effect of the independent variable (Paternal Sensitivity) on the dependent variable, controlling for the mediator (SBSK); BootLLCI = Bootstrapping lower limit confidence interval; BootULCI = Bootstrapping upper limit confidence interval; SE = Standard Error; BMI = Body Mass Index. Covariates are Child Gender, Child Ethnicity, Maternal Education, Income-to-Needs Ratio, and Full Scale IQ in 4<sup>th</sup> Grade. \*  $p < .05$ . \*\*  $p < .01$ .

Effect sizes ( $\kappa^2$ ) were .01 for both models, corresponding to a small indirect effect size (Preacher & Kelley, 2011).

Assessing partner extreme hostility as a binary outcome (0 = endorsing no indices; 1 = endorsing any indices) yielded similar results (see Table 4). Both maternal ( $b = .20$ ,  $SE = .06$ ,  $p < .001$ ) and paternal ( $b = .20$ ,  $SE = .05$ ,  $p < .001$ ) sensitivity were significant predictors of secure base script knowledge, which in turn significantly predicted higher likelihood of partner's extreme hostility ( $b = -.48$ ,  $SE = .26$ ,  $p = .06$  and  $b = -.48$ ,  $SE = .29$ ,  $p = .09$ , respectively). Moreover, coefficients for the indirect effects in both cases were significant ( $b = -.10$ , 95% CI =  $-.24, -.01$  and  $b = -.10$ , 95% CI =  $-.23, -.01$ , respectively).

To assess whether each parental sensitivity variable had a unique effect on partner's extreme hostility, we conducted mediation analyses with one parental sensitivity variable serving as the independent variable and the other as a covariate. Results showed that maternal and paternal sensitivity each contributed uniquely to the explained variance in partner's extreme hostility, as indicated in both mediation analyses remaining significant (see Table 5).

### Depressive symptoms

We next assessed whether secure base script knowledge mediated the link between parental sensitivity across the first 15 years of life and self-reported depressive symptoms at age 26. For completeness, we also conducted a mediational analysis examining whether secure base script knowledge accounted for the association between paternal sensitivity and depressive symptoms despite the focal correlation being nonsignificant ( $r = -.08$ ,  $p = .09$ ). Supporting our mediation hypothesis, results indicated that both

**Table 4.** Path coefficients and indirect effects for binary logistic regression mediation models with parental sensitivity as independent variable, secure base script knowledge as mediator, and Partner Extreme Hostility as a dependent variable.

Independent Variable ( <i>n</i> )	Path A			Path B			Path C'			Indirect Effects		
	<i>b</i>	<i>SE</i>	<i>t</i>	<i>b</i>	<i>SE</i>	<i>z</i>	<i>b</i>	<i>SE</i>	<i>z</i>	<i>Effect</i>	<i>BootLLCI</i>	<i>BootULCI</i>
Maternal Sensitivity (390)	.20**	.06	3.30	-.48 <sup>†</sup>	.26	-1.86	-.28	.27	-1.05	-.10	-.24	-.01
Paternal Sensitivity (348)	.20**	.05	4.09	-.48 <sup>†</sup>	.29	-2.22	-.27	.19	-1.37	-.10	-.23	-.01

Path A = Direct effect of the independent variable (Parental Sensitivity) on the mediator (Secure Base Script Knowledge; SBSK); Path B = Direct effect of the mediator (SBSK) on the dependent variable (Partner Extreme Hostility); Path C' = Direct effect of the independent variable (Parental Sensitivity) on the dependent variable, controlling for the mediator (SBSK); BootLLCI = Bootstrapping lower limit confidence interval; BootULCI = Bootstrapping upper limit confidence interval; SE = Standard Error. <sup>†</sup> $p < .10$  \*  $p < .05$ . \*\*  $p < .01$ .

maternal ( $b = .18$ ,  $SE = .05$ ,  $p < .001$ ) and paternal ( $b = .21$ ,  $SE = .04$ ,  $p < .001$ ) sensitivity were significant predictors of secure base script knowledge, which in turn significantly predicted depressive symptoms ( $b = -.07$ ,  $SE = .03$   $p = .007$  and  $b = -.07$ ,  $SE = .03$   $p = .02$ , respectively). The indirect coefficients were significant, as evidenced by the absence of a zero value within the bias-corrected bootstrapped confidence intervals ( $b = -.01$ , 95% CI =  $-.03$ ,  $-.001$  for both models). Approximately 4% ( $R^2 = .04$ ) of the variance in depressive symptoms was accounted for by the predictors in both models. In both models, indirect-path effect sizes were small,  $\kappa^2 = .02$ .

We also assessed whether maternal and paternal sensitivity each uniquely predicted depressive symptoms by conducting the mediation analysis with the opposite parental sensitivity variable as a covariate. In both cases the mediation model remained significant (see Table 4), indicating the unique contribution of each parental care over the first 15 years of life on depressive symptoms at age 26.

### BMI

Finally, we tested whether secure base script knowledge mediated the link between parental sensitivity across the first 15 years of life and BMI at age 26 years. Results indicated that both maternal ( $b = .18$ ,  $SE = .05$ ,  $p < .001$ ) and paternal ( $b = .21$ ,  $SE = .04$ ,  $p < .001$ ) sensitivity were significant predictors of secure base script knowledge. However, we did not find that secure base script knowledge mediated the link between parental sensitivity and BMI for either parent ( $b = -.15$ ,  $SE = .25$ ,  $p = .56$  for maternal sensitivity as an independent variable and  $b = -.05$ ,  $SE = .28$   $p = .85$  for paternal sensitivity as an independent variable).

**Table 5.** Path coefficients and indirect effects for mediation models with parental sensitivity as independent variable and secure base script knowledge as mediator, controlling for sensitivity of other parent.

Outcome ( <i>n</i> )	Path A			Path B			Path C'			Indirect Effects		
	<i>b</i>	<i>SE</i>	<i>t</i>	<i>b</i>	<i>SE</i>	<i>t</i>	<i>b</i>	<i>SE</i>	<i>t</i>	<i>Effect</i>	<i>BootLLCI</i>	<i>BootULCI</i>
<b>Maternal Sensitivity<sup>1</sup></b>												
Partner Extreme Hostility (357)	.23**	.05	4.20	-.03*	.01	-2.41	-.20	.01	-1.92	-.01	-.01	-.001
Depressive Symptoms (511)	.21**	.05	4.43	-.06*	.03	-2.22	-.20	.03	-.71	-.01	-.03	-.002
<b>Paternal Sensitivity<sup>1</sup></b>												
Partner Extreme Hostility (357)	.20**	.05	3.90	-.03*	.01	-2.41	-.01	.01	-.95	-.01	-.01	-.001
Depressive Symptoms (511)	.20**	.04	4.43	-.06*	.03	-2.22	-.02	.03	-.64	-.01	-.03	-.002

<sup>1</sup>Independent variable. Path A = Direct effect of the independent variable (Parental Sensitivity) on the mediator (Secure Base Script Knowledge; SBSK); Path B = Direct effect of the mediator (SBSK) on the dependent variable; Path C' = Direct effect of the independent variable (Parental Sensitivity) on the dependent variable, controlling for the mediator (SBSK); BootLLCI = Bootstrapping lower limit confidence interval; BootULCI = Bootstrapping upper limit confidence interval; SE = Standard Error. \*  $p < .05$ . \*\*  $p < .01$ .

## Discussion

The primary goals of this study were (1) to assess whether secure base script knowledge predicts later adult functioning, as reflected in romantic relationship quality, emotional health (i.e. depressive symptoms), and physical health (i.e. BMI); and (2) to determine the degree to which secure base script knowledge mediated links between early maternal and paternal sensitivity and these adult outcomes. Results suggested that more access to, and elaborated knowledge of the secure base script at age 18 predicted better romantic functioning quality as evidenced by lower frequencies of perceived romantic partner's extreme hostility, as well as better emotional health (i.e. fewer depressive symptoms) and physical health (i.e., lower BMI) at age 26 years. Moreover, secure base script knowledge partially mediated the link between early maternal and paternal sensitivity and later perceived romantic partner extreme hostility and depressive symptoms, although the explained amount of variance by the outcomes was modest (i.e. 4–5%). Individuals who experienced higher levels of maternal and paternal sensitivity showed more access to, and elaborated knowledge of the secure base script in late adolescence, which, in turn, was associated with less perceived romantic partner extreme hostility and fewer depressive symptoms in adulthood.

### Secure base script knowledge and romantic functioning

This study provides a partial replication and extension of previous results reported by Roisman et al. (2001). Paralleling their results, we found that attachment representations (i.e., secure base script knowledge as assessed via the ASA) mediated the association between quality of early child-parent interactions, and one aspect of romantic relationship behaviors in young adulthood (i.e. partner's extreme hostility). As mentioned, one should consider the low base rate of partner's extreme hostility when interpreting the findings; accordingly, the results may be generalized to only a small segment of the population, where extreme hostility behaviors are present. We also extended prior findings by showing that this mediational path was significant for each parent (i.e. both maternal and paternal sensitivity scores), and that each parent's sensitivity uniquely contributed to explained variance in outcomes above and beyond the other. Together, these results are consistent with an emerging literature suggesting that: (a) attachment representations in the form of secure base script knowledge are shaped by early experiences with primary caregivers (as previously reported in Steele et al., 2014; see also Nivison et al., 2020; Schoenmaker et al., 2015; Vaughn et al., 2016; Waters et al., 2017), and that (b) these representations, in turn, influence the processing of interpersonal information and behaviors in subsequent interpersonal relationships (e.g. Collins & Sroufe, 1999; Dykas & Cassidy, 2011; Waters et al., 2013, 2018).

We did not, however, find evidence in the SECCYD data to support the hypothesis that secure base script knowledge was associated with two other aspects of perceived romantic relationship quality (i.e., moderate hostility and warmth). Worth noting in this context is that Roisman et al. (2001) focused on direct observations of the quality of young adults' romantic relationships, whereas this study was based on self-reports. For this reason, it falls to future research to determine whether and how much attachment representations are predictive of a range of quality indicators in adults' romantic relationships.

## Secure base script knowledge and depressive symptoms

With regard to emotional health, we found that maternal, but not paternal sensitivity, was significantly and negatively associated with depressive symptoms, and that links between parental sensitivity and depressive symptoms were partially mediated by secure base script knowledge. Of note, the indirect effect of paternal sensitivity on depressive symptoms via secure base script knowledge proved significant in the absence of a direct effect of paternal sensitivity on depressive symptoms ( $r = -.08, p = .09$ ). Taken together, these findings partially support a core premise of attachment theory; that is, that the quality of early care, reflected in parents' provision of sensitive care, shapes the development and consolidation of later constructed mental representations, which, in turn, influence mental health (Bowlby, 1973, 1980; Sroufe et al., 2005).

Secure base script knowledge reflects individuals' expectations regarding the sequence of events following distress (i.e. seeking comfort, being effectively soothed, returning to exploration; Waters & Roisman, 2019). The failure to establish a belief in the availability of caregivers at times of need (e.g. a parent who is unresponsive, rejecting, or overinvolved), which is often a derivative of parental insensitivity (Biringen, 2000), is likely to stand in the way of constructing more access to, and elaborated knowledge of the secure base script. A schematic representation that lacks the expectation that one may be effectively soothed by close others at times of need contributes, in turn, to a reappraisal of stressful events as disproportional to one's ability to cope with them (Rueger et al., 2016; Stice et al., 2004). Ultimately, such a gap between presence of psychosocial stressors and perceived lack of support leads to more depressive symptomatology (Eberhart & Hammen, 2009; Hammen, 2005; Rudolph et al., 2000).

## Secure base script knowledge and physical health

Similar to findings reported by Farrell et al. (2019), our analyses suggested that higher levels of parental sensitivity experienced during childhood predicted lower BMI in young adulthood. Early negative parent-child relationships (e.g. non-supportive parenting, low levels of maternal warmth and nurturance) can lead to chronic distress that is associated with physiological wear and tear (Brody et al., 2015; Chen et al., 2011; Miller et al., 2011). Over time, these negative experiences may lead to impairment in inhibitory control and poor health-related behaviors, contributing to obesity risk in adulthood (Anderson & Keim, 2016). Future research that incorporates assessments of stress-regulation and biological markers of stress and metabolic function would help clarify these mechanisms.

However, in contrast to the results reported by Farrell et al. (2019), we did not find that secure base script knowledge mediated the association between early parental sensitivity and weight status as indicated by BMI. These results are somewhat surprising, given that variability in secure base script knowledge was previously found to be related to parental sensitivity during childhood (Steele et al., 2014; Waters et al., 2017), and to cardiometabolic risk, of which BMI is major contributor (Farrell et al., 2019). Our failure to replicate the mediated pathway between early parental sensitivity and BMI in adulthood found in Farrell et al. (2019) may be due to one or a combination of three major methodological differences between the two studies.

First, in Farrell et al. (2019) BMI was examined as one indicator in an overall cardiometabolic risk index, together with three other health markers (i.e. blood pressure, waist-to-hip ratio, and C-reactive protein). Second, the difference in secure base script knowledge measurements used in Farrell et al. (2019;  $AAI_{sb}$ ) versus the one used in the current study (ASA) may contribute to the different results between the studies regarding its mediating effects. Third, secure base script knowledge in Farrell et al.'s study (Farrell et al., 2019) was derived by (a) averaging assessments at two time points (versus a single time point in our study), and (b) assessing at later ages (ages 19 and 26 years, compared with age 18 in the current study).

### Study limitations and future research

Although findings of this study partially replicate and extend previous research on the role of secure base script knowledge in shaping adult functioning, two limitations should be noted. First, all outcomes in the current study were assessed via self-reports from a single follow-up assessment at around age 26 years. New research should expand our results by using multiple informants to evaluate the functioning of the target individuals (e.g., assessing romantic partners' evaluations of the target's warmth and hostility in romantic relationships). Second, as indicated by our attrition analyses, the subsample assessed in the current study was more affluent than the participants in original SECCYD NICHD sample who were lost due to attrition. This limits the generalizability of the current results. In the future, results should be replicated in more at-risk samples to enhance the external validity of this study. Third, due to the single time point administration of the ASA, we were not able to infer the stability of attachment-related information processing in early adulthood. Future research would benefit from multiple assessments of secure base script knowledge over time to trace the trajectory of growth/change in these representations. Finally, whereas the composite scores of parental sensitivity over multiple observations at different age points provided a robust representation of the quality of early child–parent interactions, it leaves open the question of whether or not sensitivity at a specific developmental period has unique and critical implications for future psychological and relational adaptations.

### Conclusion

This study provides new evidence to highlight the pathway by which early life experiences with one's primary caregivers endures to influence multiple domains of functioning in young adulthood. With recent calls for better understanding of the mechanisms linking close relationships in early life to both interpersonal and health outcomes later in life (Farrell & Stanton, 2019; Thoits, 2011), this report provides a further step towards understanding such developmental pathways via secure base script knowledge measured using the ASA. Although we partially replicated and extended some of the results from previous studies assessing the role of secure base script knowledge in the association between early parental sensitivity and later adult outcomes, additional research using multiple attachment representation measures in more diverse samples is clearly needed to determine the robustness of these mediational paths.

## Author note

A cooperative agreement (5 U10 HD027040) between the study investigators that included Cathryn Booth-LaForce, Deborah Lowe Vandell, and Glenn I. Roisman and the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development (NICHD) supported the design and data collection of the NICHD Study of Early Child Care and Youth Development (SECCYD) from birth through age 15 years. The age 18-year follow-up data collection was supported by the NICHD under Award Number R01 HD054822 to Cathryn Booth-LaForce and a grant from the NICHD under Award Number R01 HD069442 to Glenn I. Roisman supported the coding of the Attachment Script Assessments. The Charles Stewart Mott Foundation supported data collection at age 26 years under Award Number G-2017-00786 to Deborah Lowe Vandell. Current assessments of the SECCYD with a focus on adult health are supported by the National Heart, Lung, and Blood Institute under Award Number R01 HL130103 to Maria Bleil and by the NICHD under Award Number R01 HD091132 to Maria Bleil and Glenn I. Roisman. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

## Disclosure statement

In accordance with Taylor & Francis policy and my ethical obligation as a researcher, I am reporting that the authors of this manuscript have no interests to disclose.

## ORCID

Or Dagan  <http://orcid.org/0000-0002-4674-5425>

Brian E. Vaughn  <http://orcid.org/0000-0002-0035-2976>

## References

- Ainsworth, M. D. S., Blehar, M. C., Waters, E., & Wall, S. (1978). *Patterns of attachment*. Erlbaum.
- Anderson, S. E., Gooze, R. A., Lemeshow, S., & Whitaker, R. C. (2012). Quality of early maternal-child relationship and risk of adolescent obesity. *Pediatrics*, *129*(1), 132–140. <https://doi.org/10.1542/peds.2011-0972>
- Anderson, S. E., & Keim, S. A. (2016). Parent-child interaction, self-regulation, and obesity prevention in early childhood. *Current Obesity Reports*, *5*(2), 192–200. <https://doi.org/10.1007/s13679-016-0208-9>
- Bernard, K., Hostinar, C. E., & Dozier, M. (2018). Longitudinal associations between attachment quality in infancy, C-reactive protein in early childhood, and BMI in middle childhood: Preliminary evidence from a CPS-referred sample. *Attachment & Human Development*, *21*(1), 1–18. <https://doi.org/10.1080/14616734.2018.1541513>
- Biringen, Z. (2000). Emotional availability: Conceptualization and research findings. *American Journal of Orthopsychiatry*, *70*(1), 104–114. <https://doi.org/10.1037/h0087711>
- Booth-LaForce, C., Groh, A. M., Burchinal, M. R., Roisman, G. I., Owen, M. T., & Cox, M. J. (2014). V. Caregiving and contextual sources of continuity and change in attachment security from infancy to late adolescence. *Monographs of the Society for Research in Child Development*, *79*(3), 67–84. <https://doi.org/10.1111/mono.12114>
- Bowlby, J. (1969). *Attachment and loss: Vol. 1. Attachment*. Basic Books.
- Bowlby, J. (1973). *Attachment and loss: Vol. 2. Separation: Anxiety and anger*. Basic Books.
- Bowlby, J. (1980). *Attachment and loss: Vol. 3. Loss: Sadness and depression*. Basic Books.
- Bretherton, I. (1985). Attachment theory: Retrospect and prospect. *Monographs of the Society for Research in Child Development*, *50*(1–2), 3–35. <https://doi.org/http://dx.doi.org.libproxy.newschool.edu/10.2307/3333824>

- Bretherton, I. (1987). New perspectives on attachment relations: Security, communication, and internal working models. In J. Osofsky (Ed.), *Handbook of infant development* (pp. 1061–1100). Wiley.
- Bretherton, I., & Munholland, K. A. (2008). Internal working models in attachment relationships: Elaborating a central construct in attachment theory. In J. Cassidy & P. R. Shaver (Eds.), *Handbook of attachment: Theory, research and clinical applications* (2nd ed., pp. 102–127). Guilford Press.
- Brody, G. H., Yu, T., Beach, S. R. H., & Philibert, R. A. (2015). Prevention effects ameliorate the prospective association between nonsupportive parenting and diminished telomere length. *Prevention Science, 16*(2), 171–180. <https://doi.org/10.1007/s11121-014-0474-2>
- Carroll, J. E., Gruenewald, T. L., Taylor, S. E., Janicki-Deverts, D., Matthews, K. A., & Seeman, T. E. (2013). Childhood abuse, parental warmth, and adult multisystem biological risk in the coronary artery risk development in young adults study. *Proceedings of the National Academy of Sciences of the United States of America, 110*(42), 17149–17153. <https://doi.org/10.1073/pnas.1315458110>
- Chen, E., Miller, G. E., Kobor, M. S., & Cole, S. W. (2011). Maternal warmth buffers the effects of low early-life socioeconomic status on pro-inflammatory signaling in adulthood. *Molecular Psychiatry, 16*(7), 729–737. <https://doi.org/10.1038/mp.2010.53>
- Cohen, J. (1992). A power primer. *Psychological Bulletin, 112*(1), 155–159. <https://doi.org/10.1037/0033-2909.112.1.155>
- Collins, W. A., & Sroufe, L. A. (1999). Capacity for intimate relationships: A developmental construction. In W. Furman, B. Bradford Brown, & C. Feiring (Eds.), *The development of romantic relationships in adolescence* (pp. 125–147). Cambridge University Press. <https://doi.org/10.1017/CBO9781316182185.007>
- Conger, R. D. (1989). *Behavioral Affect Rating Scale (BARS)*. Developed from diverse sources for the Iowa Youth & Family. Iowa State University.
- Conger, R. D., Cui, M., Bryant, C. M., & Elder, G. H. (2000). Competence in early adult romantic relationships: A developmental perspective on family influences. *Journal of Personality and Social Psychology, 79*(2), 224–237. <https://doi.org/10.1037/0022-3514.79.2.224>
- Cosco, T. D., Prina, M., Stubbs, B., & Wu, Y.-T. (2017). Reliability and validity of the center for epidemiologic studies depression scale in a population-based cohort of middle-aged U.S. adults. *Journal of Nursing Measurement, 25*(3), 476–2017. <https://doi.org/10.1891/1061-3749.25.3.476>
- Crowell, J. A., Treboux, D., Gao, Y., Fyffe, C., Pan, H., & Waters, E. (2002). Assessing secure base behavior in adulthood: Development of a measure, links to adult attachment representations, and relations to couples' communication and reports of relationships. *Developmental Psychology, 38*(5), 679–693. <https://doi.org/10.1037/0012-1649.38.5.679>
- Dagan, O., Asok, A., Steele, H., Steele, M., & Bernard, K. (2018a). Attachment security moderates the link between adverse childhood experiences and cellular aging. *Development and Psychopathology, 30*(4), 1211–1223. <https://doi.org/10.1017/S0954579417001705>
- Dagan, O., Facompré, C. R., & Bernard, K. (2018b). Adult attachment representations and depressive symptoms: A meta-analysis. *Journal of Affective Disorders, 236*, 274–290. <https://doi.org/10.1016/j.jad.2018.04.091>
- Dagan, O., Facompré, C. R., Nivison, M. D., Roisman, G. I., & Bernard, K. (2020). Preoccupied and dismissing attachment representations are differentially associated with anxiety in adolescence and adulthood: A meta-analysis. *Clinical Psychological Science, 8*(4), 614–640. <https://doi.org/10.1177/2167702620917455>
- Davis, C. R., Usher, N., Dearing, E., Barkai, A. R., Crowell-Doom, C., Neupert, S. D., ... Crowell, J. A. (2014). Attachment and the metabolic syndrome in midlife: The role of interview-based discourse patterns. *Psychosomatic Medicine, 76*(8), 611–621. <https://doi.org/10.1097/psy.000000000000107>
- De Winter, S., Bosmans, G., & Salemink, E. (2017). Exploring the causal effect of interpretation bias on attachment expectations. *Child Development, 88*(1), 131–140. <https://doi.org/10.1111/cdev.12587>

- Diener, M. J., Geenen, R., Koelen, J. A., Aarts, F., Gerdes, V. E. A., Brandjes, D. P. M., & Hinnen, C. (2016). The significance of attachment quality for obesity: A meta-analytic review. *Canadian Journal of Behavioural Science, 48*(4), 255–265. <https://doi.org/10.1037/cbs0000050>
- Duggal, S., Carlson, E. A., Sroufe, L. A., & Egeland, B. (2001). Depressive symptomatology in childhood and adolescence. *Development and Psychopathology, 13*(1), 143–164. <https://doi.org/http://dx.doi.10.1017/S0954579401001109>
- Dykas, M. J., & Cassidy, J. (2011). Attachment and the processing of social information across the life span: Theory and evidence. *Psychological Bulletin, 137*(1), 19–46. <https://doi.org/10.1037/a0021367>
- Dykas, M. J., Woodhouse, S. S., Cassidy, J., & Waters, H. S. (2006). Narrative assessment of attachment representations: Links between secure base scripts and adolescent attachment. *Attachment & Human Development, 8*(3), 221–240. <https://doi.org/10.1080/14616730600856099>
- Eberhart, N. K., & Hammen, C. L. (2009). Interpersonal predictors of stress generation. *Personality and Social Psychology Bulletin, 35*(5), 544–556. <https://doi.org/10.1177/0146167208329857>
- Ehrlich, K., & Cassidy, J. (2019). Attachment and physical health: Introduction to the special issue. *Attachment & Human Development, 21*(1), 1–4. <https://doi.org/10.1080/14616734.2018.1541512>
- Englund, M. M., Kuo, S. I. C., Puig, J., & Collins, W. A. (2011). Early roots of adult competence: The significance of close relationships from infancy to early adulthood. *International Journal of Behavioral Development, 35*(6), 490–496. <https://doi.org/10.1177/0165025411422994>
- Farrell, A. K., & Stanton, S. C. E. (2019). Toward a mechanistic understanding of links between close relationships and physical health. *Current Directions in Psychological Science, 28*(5), 483–489. <https://doi.org/10.1177/0963721419855657>
- Farrell, A. K., Waters, T. E. A., Young, E. S., Englund, M. M., Carlson, E. E., Roisman, G. I., & Simpson, J. A. (2019). Early maternal sensitivity, attachment security in young adulthood, and cardiometabolic risk at midlife. *Attachment & Human Development, 21*(1), 70–86. <https://doi.org/10.1080/14616734.2018.1541517>
- Fraley, R. C., Roisman, G. I., Booth-LaForce, C., Owen, M. T., & Holland, A. S. (2013). Interpersonal and genetic origins of adult attachment styles: A longitudinal study from infancy to early adulthood. *Journal of Personality and Social Psychology, 104*(5), 817–838. <https://doi.org/10.1037/a0031435>
- Halligan, S. L., Herbert, J., Goodyer, I., & Murray, L. (2007). Disturbances in morning cortisol secretion in association with maternal postnatal depression predict subsequent depressive symptomatology in adolescents. *Biological Psychiatry, 62*(1), 40–46. <https://doi.org/10.1016/j.biopsych.2006.09.011>
- Hammen, C. (2005). Stress and depression. *Annual Review of Clinical Psychology, 1*, 293–319. <https://doi.org/10.1146/annurev.clinpsy.1.102803.143938>
- Hayes, A. F. (2017). *Introduction to mediation, moderation, and conditional process analysis: A regression-based approach* (2nd ed.). Guilford Press.
- Holland, A. S., & Roisman, G. I. (2010). Adult attachment security and young adults' dating relationships over time: Self-reported, observational, and physiological evidence. *Developmental Psychology, 46*(2), 552–557. <https://doi.org/10.1037/a0018542>
- Lambert, E. A., & Lambert, G. W. (2011). Stress and its role in sympathetic nervous system activation in hypertension and the metabolic syndrome. *Current Hypertension Reports, 13*(3), 244–248. <https://doi.org/10.1007/s11906-011-0186-y>
- Main, M., Goldwyn, R., & Hesse, E. (2003–2008). Adult attachment scoring and classification system. Unpublished manuscript, University of California at Berkeley.
- Marvin, R. S., & Britner, P. A. (2008). Normative development: The ontogeny of attachment. In J. Cassidy & P. R. Shaver (Eds.), *Handbook of attachment: Theory, research, and clinical applications* (2nd ed., pp. 269–294). Guilford Press.
- Miller, G. E., Chen, E., & Parker, K. J. (2011). Psychological stress in childhood and susceptibility to the chronic diseases of aging: Moving toward a model of behavioral and biological mechanisms. *Psychological Bulletin, 137*(6), 959–997. <https://doi.org/10.1037/a0024768>

- Miller, K. F., Arbel, R., Shapiro, L. S., Han, S. C., & Margolin, G. (2018). Does the cortisol awakening response link childhood adversity to adult BMI? *Health Psychology, 37*(6), 526–529. <https://doi.org/10.1037/hea0000601>
- Monteiro, L., Veríssimo, M., Vaughn, B. E., Santos, A. J., & Bost, K. K. (2008). Secure base representations for both fathers and mothers predict children's secure base behavior in a sample of Portuguese families. *Attachment & Human Development, 10*(2), 189–206. <https://doi.org/10.1080/14616730802113711>
- Nivison, M. D., Facompré, C. R., Raby, K. L., Simpson, J. A., Roisman, G. I., & Waters, T. E. A. (2020). Childhood abuse and neglect are prospectively associated with scripted attachment representations in young adulthood. *Development and Psychopathology, 1*–13. <https://doi.org/doi:10.1017/S0954579420000528>
- Preacher, K. J., & Hayes, A. F. (2008). Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behavior Research Methods, 40*(3), 879–891. <https://doi.org/10.3758/BRM.40.3.879>
- Preacher, K. J., & Kelley, K. (2011). Effect size measures for mediation models: Quantitative strategies for communicating indirect effects. *Psychological Methods, 16*(2), 93. <https://doi.org/http://dx.doi.10.1037/a0022658>
- Puig, J., Englund, M. M., Simpson, J. A., & Collins, W. A. (2013). Predicting adult physical illness from infant attachment: A prospective longitudinal study. *Health Psychology, 32*(4), 409–417. <https://doi.org/10.1037/a0028889>
- Raby, K. L., Lawler, J. M., Shlafer, R. J., Hesemeyer, P. S., Collins, W. A., & Sroufe, L. A. (2015). The interpersonal antecedents of supportive parenting: A prospective, longitudinal study from infancy to adulthood. *Developmental Psychology, 51*(1), 115. <https://doi.org/10.1037/a0038336>
- Raby, K. L., Roisman, G. I., Simpson, J. A., Collins, W. A., & Steele, R. D. (2015). Greater maternal insensitivity in childhood predicts greater electrodermal reactivity during conflict discussions with romantic partners in adulthood. *Psychological Science, 26*(3), 348–353. <https://doi.org/10.1177/0956797614563340>
- Radloff, L. S. (1977). The CES-D scale: A self-report depression scale for research in the general population. *Applied Psychological Measurement, 1*(3), 385–401. <https://doi.org/10.1177/014662167700100306>
- Rodriguez, A. C. I., Epel, E. S., White, M. L., Standen, E. C., Seckl, J. R., & Tomiyama, A. J. (2015). Hypothalamic-pituitary-adrenal axis dysregulation and cortisol activity in obesity: A systematic review. *Psychoneuroendocrinology, 62*, 301–318. <https://doi.org/http://dx.doi.10.1016/j.psyneuen.2015.08.014>
- Roisman, G. I. (2009). Adult attachment: Toward a rapprochement of methodological cultures. *Current Directions in Psychological Science, 18*(2), 122–126. <https://doi.org/10.1111/j.1467-8721.2009.01621.x>
- Roisman, G. I., Madsen, S. D., Hennighausen, K. H., Sroufe, A. L., & Collins, W. A. (2001). The coherence of dyadic behavior across parent-child and romantic relationships as mediated by the internalized representation of experience. *Attachment & Human Development, 3*(2), 156–172. <https://doi.org/10.1080/14616730126483>
- Rudolph, K. D., Constance, H., Burge, D., Lindberg, N., Herzberg, D., & Daley, S. E. (2000). Toward an interpersonal life-stress model of depression: The developmental context of stress generation. *Development and Psychopathology, 12*(2), 215–234. <https://doi.org/10.1017/S0954579400002066>
- Rueger, S. Y., Malecki, C. K., Pyun, Y., Aycocock, C., & Coyle, S. (2016). A meta-analytic review of the association between perceived social support and depression in childhood and adolescence. *Psychological Bulletin, 142*(10), 1017–1067. <https://doi.org/10.1037/bul0000058>
- Ruiz, S. K., Waters, T. E. A., & Yates, T. M. (2019). Children's secure base script knowledge as a mediator between early life stress and later behavior problems. *Attachment & Human Development, 1*–16. <https://doi.org/10.1080/14616734.2019.1672079>
- Schank, R., & Abelson, R. (1977). *Scripts, plans, goals, and understanding*. Wiley.
- Schoenmaker, C., Juffer, F., van IJzendoorn, M. H., Linting, M., van der Voort, A., & Bakermans-Kranenburg, M. J. (2015). From maternal sensitivity in infancy to adult attachment

- representations: A longitudinal adoption study with secure base scripts. *Attachment & Human Development*, 17(3), 241–256. <https://doi.org/10.1080/14616734.2015.1037315>
- Simmonds, M., Llewellyn, A., Owen, C. G., & Woolacott, N. (2016). Predicting adult obesity from childhood obesity: a systematic review and meta-analysis. *Obesity Reviews*, 17(2), 95–107.
- Skalicky, J., Vladimira, M., Roman, K., Milan, M., Tomas, R., & Vladimir, P. (2008). Evaluation of oxidative stress and inflammation in obese adults with metabolic syndrome. *Clinical Chemistry and Laboratory Medicine*, 46(4), 499–505. <https://doi.org/10.1515/CCLM.2008.096>
- Sroufe, L. A., Egeland, B., Carlson, E., & Collins, W. A. (2005). *The development of the person: The Minnesota study of risk and adaptation from birth to adulthood*. Guilford.
- Steele, R. D., Waters, T. E. A., Bost, K. K., Vaughn, B. E., Truitt, W., Waters, H. S., ... Roisman, G. I. (2014). Caregiving antecedents of secure base script knowledge: A comparative analysis of young adult attachment representations. *Developmental Psychology*, 50(11), 2526–2538. <https://doi.org/10.1037/a0037992>
- Stice, E., Ragan, J., & Randall, P. (2004). Prospective relations between social Support and depression: Differential direction of effects for parent and peer support? *Journal of Abnormal Psychology*, 113(1), 155–159. <https://doi.org/10.1037/0021-843X.113.1.155>
- Stovall-McClough, K. C., & Dozier, M. (2016). Attachment states of mind and psychopathology in adulthood. In J. Cassidy & P. R. Shaver (Eds.), *Handbook of attachment: Theory, research and clinical applications* (3rd ed., pp. 715–738). Guilford Press.
- Taylor, Z. E., Larsen-Rife, D., Conger, R. D., & Widaman, K. F. (2012). Familism, interparental conflict, and parenting in Mexican-origin families: A cultural–contextual framework. *Journal of Marriage and Family*, 74(2), 312–327. <https://doi.org/doi:10.1111/j.1741-3737.2012.00958.x>
- Thoits, P. A. (2011). Mechanisms Linking Social Ties and Support to Physical and Mental Health. *Journal of Health and Social Behavior*, 52(2), 145–161. <https://doi.org/10.1177/0022146510395592>
- Vaughn, B. E., Coppola, G., Verissimo, M., Monteiro, L., Santos, A. J., Posada, G., Korth, B. (2007). The quality of maternal secure-base scripts predicts children's secure-base behavior at home in three sociocultural groups. *International Journal of Behavioral Development*, 31(1), 65–76. <https://doi.org/10.1177/0165025407073574>
- Vaughn, B. E., Verissimo, M., Coppola, G., Bost, K. K., Shin, N., McBride, B., Krzysik, L., & Korth, B. (2006). Maternal attachment script representations: Longitudinal stability and associations with stylistic features of maternal narratives. *Attachment & Human Development*, 8(3), 199–208. <https://doi.org/10.1080/14616730600856024>
- Vaughn, B. E., Waters, T. E. A., Steele, R. D., Roisman, G. I., Bost, K. K., Truitt, W., ... Booth-Laforce, C. (2016). Multiple domains of parental secure base support during childhood and adolescence contribute to adolescents' representations of attachment as a secure base script. *Attachment & Human Development*, 18(4), 317–336. <https://doi.org/10.1080/14616734.2016.1162180>
- Waters, H. S., & Waters, E. (2006). The attachment working models concept: Among other things, we build script-like representations of secure base experiences. *Attachment & Human Development*, 8(3), 185–197. <https://doi.org/10.1080/14616730600856016>
- Waters, T. E. A., & Facompré, C. R. (in press). Measuring secure base content in the Adult Attachment Interview. In E. Waters, B. E. Vaughn, & H. S. Waters (Eds.), *Measuring attachment*. Guilford Press.
- Waters, T. E. A., Bosmans, G., Vandevivere, E., Dujardin, A., & Waters, H. S. (2015). Secure base representations in middle childhood across two Western cultures: Associations with parental attachment representations and maternal reports of behavior problems. *Developmental Psychology*, 51(8), 1013–1025. <https://doi.org/10.1037/a0039375>
- Waters, T. E. A., Brockmeyer, S. L., & Crowell, J. A. (2013). AAI coherence predicts caregiving and care seeking behavior: Secure base script knowledge helps explain why. *Attachment & Human Development*, 15(3), 316–331. <https://doi.org/10.1080/14616734.2013.782657>
- Waters, T. E. A., Facompré, C., Dagan, O., Martin, J., Johnson, W. F., Young, E. S., Shankman, J., Lee, Y., Simpson, J. A., & Roisman, G. I. (2020). Convergent validity and stability of secure base script knowledge from late adolescence to midlife. *Attachment & Human Development*.
- Waters, T. E. A., Fraley, R. C., Groh, A. M., Steele, R. D., Vaughn, B. E., Bost, K. K., ... Roisman, G. I. (2015). The latent structure of secure base script knowledge. *Developmental Psychology*, 51(6), 823–830. <https://doi.org/10.1177/0963721414541462.Self-Control>

- Waters, T. E. A., Raby, K. L., Ruiz, S. K., Martin, J., & Roisman, G. I. (2018). Adult attachment representations and the quality of romantic and parent–child relationships: An examination of the contributions of coherence of discourse and secure base script knowledge. *Developmental Psychology, 54*(12), 2371–2381. <https://doi.org/10.1037/dev0000607>
- Waters, T. E. A., & Roisman, G. I. (2019). The secure base script concept: An overview. *Current Opinion in Psychology, 25*, 162–166. <https://doi.org/10.1016/j.copsyc.2018.08.002>
- Waters, T. E. A., Ruiz, S. K., & Roisman, G. I. (2017). Origins of secure base script knowledge and the developmental construction of attachment representations. *Child Development, 88*(1), 198–209. <https://doi.org/10.1111/cdev.12571>
- Wechsler Abbreviated Scale of Intelligence. (1999). *The Psychological Corporation*. Harcourt Brace & Company.
- Wilson, O. W. A., Bopp, C. M., Papalia, Z., & Bopp, M. (2019). Objective vs self-report assessment of height, weight and body mass index: Relationships with adiposity, aerobic fitness and physical activity. *Clinical Obesity, 9*(5), e12331. <https://doi.org/10.1111/cob.12331>
- Zhao, X., Lynch, J. G., Jr, & Chen, Q. (2010). Reconsidering Baron and Kenny: Myths and truths about mediation analysis. *Journal of Consumer Research, 37*(2), 197–206. <https://doi.org/10.1086/651257>